

Nucleon-Nucleon Scattering under Spin-Isospin Reversal in Large- N_c QCD

Thomas D. Cohen* and Daniel C. Dakin†

Department of Physics, University of Maryland, College Park, MD 20742-4111

(Dated: March, 2003)

The spin-flavor structure of certain nucleon-nucleon scattering observables derived from the large N_c limit of QCD in the kinematical regime where time-dependent mean-field theory is valid is discussed. In previous work [1] this regime was taken to be where the external momentum was of order N_c which precluded the study of differential cross sections in elastic scattering. Here it is shown that the regime extends down to order $N_c^{1/2}$ which includes the higher end of the elastic regime. The prediction is that in the large N_c limit, observables describable via mean-field theory are unchanged when the spin and isospin of either nucleon are both flipped. This prediction is tested for proton-proton and neutron-proton elastic scattering data and found to fail badly. We argue that this failure can be traced to a lack of a clear separation of scales between momentum of order $N_c^{1/2}$ and N_c^1 when N_c is as small as three. The situation is compounded by an anomalously low particle production threshold due to approximate chiral symmetry.

PACS numbers: 11.15.Pg, 13.88.+e, 25.60.Bx

Calculating observables for nucleon-nucleon scattering directly from QCD is a formidable task. Given our experience with other hadronic observables, one might hope to make qualitative predictions by exploiting the large N_c limit of QCD; the limit in which the number of colors, N_c , becomes large and $1/N_c$ is regarded as an expansion parameter [2, 3]. Experience in the single baryon sector shows that one of the striking features of large N_c QCD is a simple spin-flavor structure associated with a contracted $SU(2N_f)$ symmetry which becomes exact as $N_c \rightarrow \infty$ [4]. There has been considerable work over the past several years on the spin-flavor structure of the nucleon-nucleon potential appropriate for low momentum ($p \sim N_c^0$) scattering. In particular, it has been established that the leading order terms in the potential are those for which the t -channel exchange has the same angular momentum and isospin [5]. Furthermore, this picture is consistent with both single-meson and multiple-meson exchange [6]. While this is of interest theoretically, it is of limited direct phenomenological consequence since the potential cannot be directly measured. Recently, ref. [1] noted that the spin-isospin dependence of certain nucleon-nucleon scattering observables was calculable up to $1/N_c$ corrections. In particular, they showed that for large N_c those observables which are calculable in principle from time-dependent mean-field theory (TDMFT) are invariant under simultaneous spin and isospin flips of either incident particle.

Before proceeding, it is necessary to clarify exactly what this restriction to observables calculable in TDMFT implies. Firstly, as originally noted by Witten [3] there is a kinematic constraint since TDMFT is inapplicable for momenta of order N_c^0 , but becomes exact at large N_c for momenta of order N_c^1 (which implies velocities of order N_c^0 since the nucleon mass goes N_c^1). Secondly, mean-field theory only becomes exact in the large N_c limit for observables without multi-particle correlations,

i.e. for observables associated with the collective flow of currents such as baryon number or energy. Examples of such observables include the differential cross section for baryon number (in the form of nucleons) to flow out from a collision (which in general produces many mesons) at a fixed angle. Since the observable integrates over all of the final meson states, it is independent of multiple particle correlation and depends only on the collective flow of baryon number. Thus it is calculable in TDMFT and subject to the spin-isospin constraint given above.

In ref. [1] it was pointed out that the most familiar nucleon-nucleon scattering observables—the total cross section, the total-elastic cross section and the differential cross section for elastic scattering—were not in the class of observables describable in mean-field theory in the kinematic regime of $p \sim N_c^1$. The total cross section includes contributions from nearly-forward scattering and forward scattering is not accessible in semiclassical treatments such as TDMFT [10]. Similarly, the total-elastic cross section is inaccessible. The differential elastic cross section in this kinematic regime is inaccessible in mean-field theory for a different reason. When $p \sim N_c$, one is well above the threshold for meson production (which scales as $p \sim N_c^{1/2}$ for an incident kinetic energy $p^2/(2M_N) \sim N_c^0$) and hence the typical collision is inelastic. The extraction of the elastic differential cross section thus depends strongly on multi-particle correlations in that it totally excludes the many-particle final states. As such it cannot be computed in TDMFT and the spin-isospin constraint does not apply.

As a practical matter, the exclusion of these more standard observables has made it very difficult to test the large N_c spin-flavor predictions of ref. [1]. We know of no nucleon-nucleon scattering experiments in the kinematic regime of $p \sim N_c$ (which we can take to be $p \sim M_N$) where the data has been reported as a differential cross section for the outgoing nucleons with the

outgoing mesons integrated over, and for which data is available for both protons and neutrons with polarized targets and/or beams.

Clearly it would be very useful to have some experimentally accessible observable that is also computable in TDMFT. Ideally, we would hope to find some observable which is readily available experimentally and of the correct class. Since the most readily available data is for the total and elastic cross sections discussed above, it would be extremely useful if there were some circumstance in which one of these may be reliably computed in TDMFT. The argument that forward scattering is not computable in TDMFT is quite generic and holds for all regimes. In contrast, the argument that the differential cross section for elastic scattering is not computable in TDMFT depended on Witten's kinematic condition that $p \sim N_c$ for TDMFT to be valid. The key theoretical point of the present paper is that Witten's condition is overly restrictive. While TDMFT is clearly inapplicable in the regime where $p \sim N_c^0$ and is clearly applicable in the regime of $p \sim N_c^1$, there is an intermediate regime of $p \sim N_c^{1/2}$. This intermediate regime includes the threshold for meson production so that one can be just below the meson production threshold and still be in this regime. Here we will argue that TDMFT is, in fact, applicable in the intermediate regime at sufficiently large N_c and as such one can have observables associated with current flows. Below meson production, however, the differential cross section for baryon flow into a given direction is simply the elastic differential cross section.

We begin by reviewing the argument that led to the spin-isospin constraint under investigation. First, let us consider nucleons in the large N_c limit. Witten argued that they could be described by a Hartree-type approximation. From this, it was deduced that the nucleon size is of order unity and that the mass is of order N_c [3]. Witten also noted that quantities in large N_c QCD scale with $1/N_c$ in the same manner that the analogous quantities in soliton models scale with g^2 , where g is the coupling constant. This suggests that nucleons can be interpreted as solitons of mesonic fields. This notion was exemplified in terms of the Skyrme model—a model in which baryons are topological solitons of a nonlinear chiral field U_0 . The large N_c limit corresponds to the classical solution [7]. For the ground state baryons the field takes the *hedgehog* form $U_0(x) = \exp[i\theta(r)\vec{\tau} \cdot \hat{r}]$, where the isospin and coordinate space vectors are coupled; the detailed functional form of the chiral angle, $\theta(r)$, is irrelevant for our purpose. The hedgehog breaks both angular momentum and isospin and thus corresponds to states of more than one spin and isospin. To get physical states with well-defined spin and isospin we need to project the hedgehog, which at large N_c can be done via a semiclassical “cranking” technique. In this formalism a set of collective coordinates associated with isospin rotations is introduced $\{a_i, a_0^2 + \vec{a} \cdot \vec{a} = 1\}_{i=0}^3$ and it is convenient to

represent them as a matrix: $A = a_0 + \vec{a} \cdot \vec{\tau} \in SU(2)$. Then the matrix $U(t) \equiv AU_0A^\dagger$ contains all the information about the collective dynamics. These dynamics can be quantized [8, 9] to yield states of definite spin (J) and isospin (I), described by wavefunctions proportional to the Wigner D functions: $\Psi(A) \propto D_{m, m_I}^{J=I}(A)$.

Now, let us now consider nucleon-nucleon scattering in light of the large N_c limit in the context of the Skyrme model. We use the Skyrme model here to make the argument concrete, but the result is expected to be completely model independent. As noted by Witten, the strength of the nucleon-nucleon interaction is of order N_c and the scattering process can be described by TDMFT. Since the interaction and mass is of order N_c , the nucleon's momentum must also be of order N_c ($p \sim N_c$) to ensure a smooth limit for scattering observables as $N_c \rightarrow \infty$. Initially, the nucleons are well separated and approach each other along the beam direction, $\hat{n}$ with an impact parameter, $\hat{b}$. In the Skyrme picture each is taken as a hedgehog, parameterized by its own A matrix and wavefunction $\Psi(A)$, with spin ($m^{(1,2)}$) and isospin ($m_I^{(1,2)}$) projections. In TDMFT, the observable must be written in terms of an operator $f(A_1, A_2)$. After quantization, this corresponds to integrating over the collective variables weighted by the appropriate collective wavefunctions. The resulting operator loses any explicit dependence on the collective variables and instead are expressed in terms of the spin or isospin. Moreover, these quantities only appear in the form $\vec{S}/\mathcal{I}$ or $\vec{I}/\mathcal{I}$, where $\mathcal{I}$ is the hedgehog Skyrminion's moment of inertia. Since $\mathcal{I}$ is of the order N_c [8], these terms may be neglected at leading order. To compute an observable, we must calculate the expectation value of the associated operator and integrate over the impact parameter and collective variables. Reference [1] implemented this procedure and arrived at the following form for an observable:

$$\begin{aligned} \sigma_{m^{(1)}, m_I^{(1)}, m^{(2)}, m_I^{(2)}}^O &= \int d^2\vec{b} dA_1 dA_2 \left| D_{m^{(1)}, m_I^{(1)}}^{1/2}(A_1) \right|^2 \\ &\times \left| D_{m^{(2)}, m_I^{(2)}}^{1/2}(A_2) \right|^2 f(A_1, A_2), \end{aligned} \quad (1)$$

where $\int dA$ is the invariant Haar measure on $SU(2)$. We note that the spin and isospin projection dependence is contained in the Wigner D functions and that the time reversal invariance of the rotation operator dictates that $\left| D_{m, n}^{1/2}(A) \right|^2 = \left| D_{-m, -n}^{1/2}(A) \right|^2$. This finally leads us to the condition that observables are unchanged at leading order in $1/N_c$ when one simultaneously flips both the spin and isospin of either nucleon. As an example, consider this condition for an observable which averages over the direction of all detected particles, thus leaving the beam direction, $\hat{n}$, and spins as the only remaining vectors. Insisting that it be invariant under time reversal and parity,

we arrive at the general form

$$\begin{aligned} \sigma^O = & A_0 + B_I(\vec{\sigma}^{(1)} \cdot \vec{\sigma}^{(2)})(\vec{\tau}^{(1)} \cdot \vec{\tau}^{(2)}) \\ & + C_I(\hat{n} \cdot \vec{\sigma}^{(1)})(\hat{n} \cdot \vec{\sigma}^{(2)})(\vec{\tau}^{(1)} \cdot \vec{\tau}^{(2)}), \end{aligned} \quad (2)$$

where A_0, B_I, C_I are functions of energy. If one averages over the spins, as we would for an unpolarized measurement, we are left with only the isoscalar A_0 . Therefore, the result will be independent of isospin.

The key issue which must be addressed is the kinematic conditions under which the preceding analysis is valid. Clearly we must assume that N_c is sufficiently large. Moreover, Witten noted that at low momenta, $p \sim N_c^0$, one does not have a smooth large N_c limit for scattering observables, while at larger momenta, $p \sim N_c$, those observables calculable in TDMFT will have smooth limits provided we keep the velocities fixed as we take N_c to infinity. The issue of relevance here is the determination of the minimum velocity for TDMFT to be valid. To focus sharply on this issue consider some quantity, Θ , which is computable in TDMFT. At large N_c , Θ is a function of the velocity (which scales as p/N_c). Consider now the limit of this function as the velocity approaches zero. The quantity involves a double limit, $N_c \rightarrow \infty$ and $v \rightarrow 0$, and the limit is not uniform. If the large N_c limit is taken first, one can smoothly take v down to zero. However, if one takes $v \rightarrow 0$ first, then for any given N_c one passes through the regime $p \sim N_c^0$ prior to reaching $v = 0$. However, the regime $p \sim N_c^0$ is outside the validity of TDMFT, and one expects Θ to have no smooth large N_c limit.

Consider the combined zero velocity and large N_c limits taken in the following way:

$$N_c^{-1} \rightarrow 0, \quad v \rightarrow 0 \quad \text{with} \quad vN_c^{1/2} \sim pN_c^{-1/2} \rightarrow \text{fixed}. \quad (3)$$

This limit is interesting in that it corresponds simultaneously to zero velocity and infinite momentum. As we shall argue below, the infinite momentum nature of the limit justifies the use of TDMFT and, thus, this combined limit is both smooth and has the same result as taking the large N_c limit first and then the zero velocity limit.

Formally, we see that as $N_c \rightarrow \infty$, velocities of order $N_c^{-1/2}$ are in the semi-classical regime and, hence, can be described by TDMFT. Moreover, the usual differential cross section in this regime and just below the meson production threshold is exactly the same as the observable associated with the differential flow of baryon number through various angles and as such is computable in TDMFT.

The argument that the limiting procedure given above leads to a quantity computable in TDMFT is straightforward. TDMFT is semi-classical in nature and the fundamental issue is under what kinematic conditions the classical behavior sets in. We know that the momentum must

be high enough so that the nucleon's deBroglie wavelength is small compared to the size of the interaction region. Also, the variation in the effective wavelength as it traverses the interaction region must remain small compared to the wavelength itself. This condition must hold even when the interaction strength becomes large, as in our case since it scales as N_c^1 . Simple potential model calculations with masses and potential strengths both scaling as N_c^1 show that the semi-classical condition is reached for $p \sim N_c^{1/2}$. With this it is highly plausible that the semi-classical approximation is valid for $p \sim N_c^{1/2}$ for large N_c QCD. Since the key issues here are spin and flavor, one condition for the semi-classical regime to be valid is that the nucleon's velocity must be rapid enough so that the characteristic rotation time of a hedgehog is long compared to the characteristic interaction time. The characteristic rotation time depends on the moment of inertia $\mathcal{I}$ and is given by $\mathcal{I}/S$ which is order N_c ; the characteristic interaction time is R/v , where $R \sim N_c^0$ is the size of the nucleon. Reference [1] noted that the condition holds for velocities of order N_c^0 which yields interaction times of order unity. Here we simply note that the condition also holds in the large N_c limit for $v \sim N_c^{-1/2}$ which yields interaction times of order $N_c^{1/2}$ which is parametrically much smaller than the rotation times.

We also note that the semi-classical approach is known to fail for forward scattering [10], meaning one cannot make valid predictions in the small angular region around the beam direction, $\hat{n}$.

Typical elastic nucleon-nucleon scattering experiments consist of a proton beam with specified polarization incident on a proton or neutron target. Detectors in the reaction chamber can record the lab angle of scattering, the energy, and the polarization of the scattered nucleon. From this information, various observables can be calculated to describe the event. Let us consider two examples: the unpolarized differential cross section ($d\sigma/d\Omega$) for elastic collisions and the polarization asymmetry in the cross section ($\mathcal{A}$).

The unpolarized differential cross section, $d\sigma/d\Omega$, is obtained by counting the number of nucleons scattered into the solid angle $d\Omega$ and averaging over the polarization of the beam. Since this is an unpolarized measurement, and, hence, independent of the nucleon spins, the spin-flavor condition of eq. (2) implies $d\sigma/d\Omega$ must be an isoscalar, meaning that np and pp scattering should be equal up to $1/N_c$ corrections. In fig. 1(a) we show $d\sigma/d\Omega$ versus the scattering angle as measured in the nucleon's center of mass frame for a proton beam with kinetic energy of $T_{lab} = 279$ MeV (which is just below the pion production threshold) using data from ref. [11]. For the reason mentioned above, we have omitted forward angles.

The polarization asymmetry, $\mathcal{A}$, quantifies the depen-




FIG. 1: (a) Elastic $d\sigma/d\Omega$ and (b) $\mathcal{A}_{pp}$ and $-\mathcal{A}_{np}$ for $T_{lab} = 279$ MeV as taken from [11]. The dashed line is for pp and the solid line is for np.

dence of the differential cross section on the beam polarization. Consider a target of protons (p) with a set spin ($m_t = +$, for example) and a beam of nucleons N (either n or p) with set spins ($m_b = \pm$). The differential cross section obtained for this arrangement will be written as $d\sigma_{m_b m_t}^N$ with the $d\Omega$ suppressed since it is irrelevant here. The asymmetry for pp ($\mathcal{A}_{pp}$) and np ($\mathcal{A}_{np}$) scattering is then written as

$$\mathcal{A}_{pp} = \frac{d\sigma_{++}^p - d\sigma_{-+}^p}{d\sigma_{++}^p + d\sigma_{-+}^p} \quad (4)$$

$$\mathcal{A}_{np} = \frac{d\sigma_{++}^n - d\sigma_{-+}^n}{d\sigma_{++}^n + d\sigma_{-+}^n}. \quad (5)$$

Under spin and isospin reversal of the beam nucleons, $d\sigma_{++}^n \rightarrow d\sigma_{-+}^p$ and $d\sigma_{-+}^n \rightarrow d\sigma_{++}^p$, meaning that eq. (2) implies that at leading order in $1/N_c$, $\mathcal{A}_{np} = -\mathcal{A}_{pp}$. In figure 1(b) we show the $\mathcal{A}$ versus the scattering angle as measured in the nucleon's center of mass frame for a proton beam with kinetic energy of $T_{lab} = 279$ MeV.

Clearly, the large N_c predictions for the two observables considered above fail quite badly. These two cases are typical of the disagreement found between the proposed spin-isospin dependence and data collected from nucleon-nucleon elastic scattering experiments in this kinematic range. Moreover, we cannot go to higher energies above 279 MeV because we will hit the pion pro-

duction threshold. Above the threshold, the elastic differential cross section is not equal to the full collective flow of baryon as baryons can flow in a given direction accompanied by meson production. Thus, above threshold the elastic differential cross section is not computable in TDMFT.

Why do the large N_c predictions for these observables fail so badly? One very plausible explanation for failure of the large N_c predictions with these kinematics is that the velocity is insufficient to bring us to the required regime. As argued before, the beam energy must be high enough to achieve the semi-classical scattering regime. However, as the beam's kinetic energy increases, it reaches the threshold for meson production. Kinematically the threshold occurs at $T_{lab} = \frac{1}{2}M_N v_t^2 = 2m_\pi$. Thus the maximum velocity before the meson production channel opens is $v_t = 2\sqrt{m_\pi/M_N} \sim N_c^{-1/2}$. As argued above, the spin-flavor arguments here depend on the interaction times being short compared to the characteristic rotation time scales. Were this not true, additional terms proportional to $\vec{S}/I$ and $\vec{I}/I$ could be added to eq. (2). In the kinematics of ref. [1] with $v \sim N_c^0$ the ratio of the interaction time to the rotation time was of order $1/N_c$ implying that corrections due to these possible extra terms were suppressed by $1/N_c$. With the present kinematics the ratio is $N_c^{-1/2}$ and the additional terms have a characteristically smaller suppression. Of course, if N_c is large enough one expects the leading order expression to become increasingly good. However, for $N_c = 3$ it is by no means clear that what amounts to an expansion in $N_c^{-1/2}$ will usefully converge.

There is an additional issue which limits the usefulness of the large N_c expansion for these observables. The threshold velocity for meson production is proportional to the square root of the meson mass. Moreover, elastic scattering is only associated with collective baryon flow and thus calculable by TDMFT when one is below the threshold for *all mesons*. The pion is the lightest meson and thus its threshold velocity sets the maximum for which the elastic differential cross section is computable in TDMFT. It is noteworthy that the pion mass is very small on hadronic scales. This is due to the pseudo-Goldstone nature of the pion and the small values of the up and down quark masses. Thus, the pion mass is small for reasons which have nothing to do with N_c . If we denote the ratio of m_π to a typical hadronic scale as β , the threshold velocity $v_t = 2\sqrt{m_\pi/M_N} \sim \beta^{1/2} N_c^{-1/2}$. Therefore, chiral effects suppress the velocity threshold by a factor of $\beta^{1/2}$ relative to the natural scales in $1/N_c$ counting. This suppression could be responsible for pushing the elastic regime below the regime for which the large N_c expansion is useful.

In summary, we have shown in principle that the spin and flavor dependence of the elastic differential cross section in non-forward directions at energies just below the

meson production threshold can be computed in large N_c QCD. In practice, the empirical scattering data is poorly described by the large N_c relations. We attribute this to the slow convergence of the expansion which is, in effect, an expansion in $(\beta N_c)^{-1/2}$, where β is the ratio of m_π to typical hadronic scales.

The support of the U.S. Department of Energy for this research under grant DE-FG02-93ER-40762 is gratefully acknowledged.

* Electronic address: cohen@physics.umd.edu

† Electronic address: dcdakin@physics.umd.edu

- [1] T.D. Cohen, B.A. Gelman, Phys. Lett. **B540**, 227 (2002).
- [2] G. 't Hooft, Nucl. Phys. **B72**, 461 (1974).
- [3] E. Witten, Nucl. Phys. **B160**, 57 (1979).
- [4] J.L. Gervais, B. Sakita, Phys. Rev. Lett. **52**, 87 (1984);
J.L. Gervais, B. Sakita, Phys. Rev. **D30**, 1795 (1984);
R.F. Dashen, A.V. Manohar, Phys. Lett. **B315**, 425 (1993);
R.F. Dashen, A.V. Manohar, Phys. Lett. **B315**, 438 (1993);
R. Dashen, E. Jenkins, A.V. Manohar, Phys. Rev. **D49**, 4713 (1994);
C. Carone, H. Georgi, S. Osofsky, Phys. Lett. **B322**, 227 (1994);
M. Luty, J. March-Russell, Nucl. Phys. **B426**, 71 (1994);
R. Dashen, E. Jenkins, A.V. Manohar, Phys. Rev. **D51**, 3697 (1995).
- [5] D.B. Kaplan, M.J. Savage, Phys. Lett. **B365**, 244 (1996);
D.B. Kaplan, A.V. Manohar, Phys. Rev. **C56**, 76 (1997);
D.O. Riska, Nucl. Phys. **A710**, 55 (2002).
- [6] M.K. Banerjee, T.D. Cohen, B.A. Gelman, Phys. Rev. **C65**, 034011 (2002);
A.V. Belitsky, T.D. Cohen, Phys. Rev. **C65**, 064008 (2002);
T.D. Cohen, Phys. Rev. **C66**, 064003 (2002).
- [7] I. Zahed, G.E. Brown, Phys. Reports **142**, 1 (1986).
- [8] G.S. Adkins, C.R. Nappi, E. Witten, Nucl. Phys. **B228**, 552 (1983).
- [9] T.D. Cohen, W. Broniowski, Phys. Rev. **D34**, 3472 (1986).
- [10] L.D. Landau, E.M. Lifshitz, *Quantum Mechanics: Non-relativistic Theory*, 3d ed. (Pergamon, Elmsford, N.Y., 1977).
- [11] M. Rentmeester. 2002. NN-Online. Nijmegen, The Netherlands: University of Nijmegen; available from <http://nn-online.sci.kun.nl>.